



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
&= \frac{2\pi a^2 b^3 c^3 \sin A}{\sqrt{(\{4a^2 c^2 - [a^2 + c^2 - b^2]^2\}^3)}} = \frac{\pi a^2 b^3 c^3 \sin A}{32\sqrt{(\{s[s-a][s-b][s-c]\}^3)}} \\
&= \frac{\pi a^2 b^2 c^2}{[4 \text{ area of triangle}]^2} = \pi R^2.
\end{aligned}$$

CALCULUS.

160. Proposed by B. F. FINKEL, A.M., M.Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

A dog at the vertex of a right conical hill pursues a fox at the foot of the hill. How far will the dog run to catch the fox, if the dog runs directly towards the fox at all times, and the fox is continually running around the hill at its foot, the velocity of the dog being 6 feet per second, the velocity of the fox being 5 feet per second, the hill being 100 feet high and 200 feet in diameter at the base?

Note by J. E. SANDERS, Hackney, Ohio.

The latter part of the solution given in the April MONTHLY is not correct. The dog cannot run towards the fox at all times and keep between the fox and the vertex of the hill. If we suppose only that he runs so that he is directly between the fox and the vertex, the following is the solution:

Let the surface of the hill be rolled out flat. Then the radius of the fox's path = $100\sqrt{2}$ feet = a . Put $n = \frac{5}{6}$, s = length of dog's path.

$$ds = [a/n]d\theta = \sqrt{r^2 + [dr/d\theta]^2}.$$

$\therefore d\theta = \frac{ndr}{\sqrt{a^2 - n^2 r^2}}$, which gives $\theta = \sin^{-1}[nr/a]$ when integrated. When the dog catches the fox $r = a$, and
 $\therefore \theta = \sin^{-1}n$ and $s = [a/n]\theta = 120\sqrt{2}\sin^{-1}[\frac{5}{6}] = 167.178$ feet.

No solution of Problem 165 has been received. See April number, 1903, page 115.

166. Proposed by T. N. HAUN, Mohawk, Tenn.

Find the volume of the solid formed by the revolution of the curve $[y^2 + x^2] = a^2[x^2 - y^2]$ round the axis of x .

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$$[y^2 + x^2]^2 = a^2[x^2 - y^2].$$

$$\therefore y^4 + 2x^2y^2 + x^4 = a^2x^2 - a^2y^2. \quad \therefore y^2 = \frac{1}{2}\{a\sqrt{a^2 + 8x^2} - a^2 - 2x^2\}.$$